

Exact Positive Icosahedral Cubature on Fixed Radial Class-I Grids

Quadratic-field interoperability, exact replay,
and typed compatibility with Papers I–V

Matt Newman

ORCID: [0009-0007-5468-3125](https://orcid.org/0009-0007-5468-3125)

23 August 2026

Preprint, Version 1.0

Status and dependency boundary. The cubature theorems are consequences of the fixed radial Class-I geometry, full icosahedral symmetry, and exact invariant moment systems over $\mathbb{Q}(\sqrt{5})$. The OMEGA quotient algebra is used only to encode coefficients already obtained from that geometry. Its 15-bit floor contract is a separate finite replay result. The Papers I–V global object supplies a typed ambient assembly and an exact coordinate-transport interface; it is not an antecedent of the cubature proof. The all-frequency statement is a conjecture. No claim of novelty, priority, peer review, or journal acceptance is made.

Abstract

Let $X_f \subset \mathbb{S}^2$ be the complete frequency- f radial Class-I icosahedral grid. Using the full reflection group H_3 , Sobolev’s invariant-cubature reduction, and exact arithmetic in $\mathbb{Q}(\sqrt{5})$, we certify strictly positive orbit-constant probability rules on X_f for $1 \leq f \leq 8$. Their exact algebraic degrees are

$$5, 9, 11, 15, 17, 21, 23, 27,$$

respectively. Exact nonzero residuals at the first omitted invariant degrees prove that these are exact degrees, not only lower bounds.

Independently, the quotient algebra

$$K_\Omega = \mathbb{Q}(T)[\lambda]/(\lambda^2 - T\lambda - T^2)$$

admits the exact identification $\sqrt{5} = 2\lambda/T - 1$. Hence every cubature coefficient $a + b\sqrt{5}$ has the two-channel representation $(a - b) + (2b/T)\lambda$. This is a genuine coefficient-field interoperability map, but it is not a source of the icosahedral theorem.

Finally, conditional on the frozen strict enclosure of the distinguished real $\lambda_\Omega = \varphi/(\varphi + \pi)$, exact integer replay verifies

$$\lfloor \lambda_\Omega d \rfloor = \left\lfloor \frac{339949771344778}{10^{15}} d \right\rfloor \quad (0 \leq d < 2^{15}).$$

A determinant-one Farey certificate proves that the first disagreement over all nonnegative integers occurs at $d = 42,189,065$. All large coefficients and proof decisions are preserved in the deterministic verification record identified in Section 10.

The global-object integration extends the multi-sorted object associated with Papers I–V by a cubature sort. Its radial branch already contains the equal-weight measure μ_f on X_f ; the present weights define a different positive, full-support measure ν_f on the same nodes. For $f \geq 2$, $\nu_f(H_6) = 0$, so orbit reweighting cancels the equal-weight degree-six obstruction rather than reinterpreting it. Exact SRSD encoding transports the rule without changing any decoded moment. This is a conservative typed interface, not a new derivation of either branch.

Contents

1	Scope and separation of claims	2
2	Fixed Class-I grids and orbit masses	3
3	Invariant reduction and exact moments	3
4	Exact positive rules for frequencies one through eight	4
4.1	Compact exact formulas for $f \leq 4$	5
4.2	Exact sign decisions	6
5	The exact OMEGA coefficient-field representation	6
6	Finite 15-bit replay and the first disagreement	7
7	The all-frequency convex formulation	8
8	Compatibility with the Papers I–V global object	9
8.1	Same nodes, distinct measures	9
8.2	Arithmetic specialization and canonical records	10
8.3	Exact transport through SRSD coordinates	10
9	Provenance and literature boundary	11
10	Reproducibility	11
11	Conclusion	12

1 Scope and separation of claims

This paper joins several exact statements at typed interfaces, not at a common mathematical origin:

Layer	Statement	Status
Radial geometry	Definition and symmetry of X_f , with $ X_f = 10f^2 + 2$	Imported from [5]
Invariant reduction	H_3 -invariant exactness reduces to invariant moments	Classical; hypotheses checked [8]
Positive cubature	Exact positive rules and first failures for $1 \leq f \leq 8$	Exact finite computation here
OMEGA algebra	$\mathbb{Q}(\sqrt{5})$ embeds by $\sqrt{5} \mapsto 2\lambda/T - 1$	Exact algebra here
Finite floor bridge	15-bit equality and first-disagreement minimality	Exact, conditional on the frozen enclosure
Papers I–V object	Adjoin the cubature measures and exact interface laws	Conservative typed extension of [7]
All frequencies	Positive rules through the proposed threshold for $f \geq 9$	Open conjecture

Deleting the OMEGA sections and files leaves every cubature theorem and proof intact. Conversely, the OMEGA algebra acquires no geometric theorem merely by representing a coefficient produced by

the cubature calculation. Likewise, deleting the SRSD coordinate-transport interface leaves the native cubature calculation unchanged.

2 Fixed Class-I grids and orbit masses

Put

$$\varphi = \frac{1 + \sqrt{5}}{2}$$

and take the icosahedral face with vertices

$$v_1 = (0, 1, \varphi), \quad v_2 = (1, \varphi, 0), \quad v_3 = (\varphi, 0, 1).$$

For integers $a, b, c \geq 0$ with $a + b + c = f$, define

$$p_{abc} = av_1 + bv_2 + cv_3, \quad x_{abc} = \frac{p_{abc}}{\|p_{abc}\|}.$$

The complete grid X_f is the full icosahedral orbit union of these barycentric nodes, with duplicates on common edges and vertices identified. The exact count

$$|X_f| = 10f^2 + 2$$

and the radial construction are imported from [5].

Let H_3 denote the full icosahedral reflection group. Its orbits on X_f have representatives

$$\Lambda_f = \{(a, b, c) : a \geq b \geq c \geq 0, a + b + c = f\}.$$

If $\mathcal{O}_{abc} = H_3 x_{abc}$, its size is 12 at a vertex, 30 at an edge midpoint, 60 at a nonmidpoint edge node, and, for an interior face node, 20, 60, or 120 according as all three barycentric entries are equal, exactly two are equal, or all are distinct. These sizes sum to $10f^2 + 2$.

Definition 2.1 (Orbit-mass rule). For total orbit masses W_{abc} , define

$$Q_f[g] = \sum_{(a,b,c) \in \Lambda_f} \frac{W_{abc}}{|\mathcal{O}_{abc}|} \sum_{x \in \mathcal{O}_{abc}} g(x).$$

Thus W_{abc} , not $W_{abc}/|\mathcal{O}_{abc}|$, is the total probability assigned to the orbit. The rule is strictly positive when every $W_{abc} > 0$, and it is probability normalized when $\sum_{\Lambda_f} W_{abc} = 1$.

Throughout, σ denotes normalized spherical area: $\int_{\mathbb{S}^2} 1 \, d\sigma = 1$.

3 Invariant reduction and exact moments

Let

$$V_{\text{ico}} = \{(0, \pm 1, \pm \varphi), (\pm 1, \pm \varphi, 0), (\pm \varphi, 0, \pm 1)\}, \quad R^2 = \varphi + 2,$$

and write $\hat{v} = v/R$. With P_n the Legendre polynomial of degree n , define on $\mathbb{S}^2$

$$H_n(x) = \sum_{v \in V_{\text{ico}}} P_n(x \cdot \hat{v}), \quad n = 6, 10.$$

The normalizations used by the certificate are

$$P_6(t) = \frac{231t^6 - 315t^4 + 105t^2 - 5}{16}$$

and

$$P_{10}(t) = \frac{46189t^{10} - 109395t^8 + 90090t^6 - 30030t^4 + 3465t^2 - 63}{256}.$$

Their homogeneous extensions are H_3 -invariant harmonic polynomials. In particular,

$$\begin{aligned} H_6(V) &= \frac{132}{25}, & H_6(E) &= -\frac{33}{20}, \\ H_{10}(V) &= \frac{988}{625}, & H_{10}(E) &= -\frac{247}{160}, \end{aligned}$$

on a fivefold vertex axis V and a twofold edge-midpoint axis E . Also

$$\int_{\mathbb{S}^2} H_6 d\sigma = \int_{\mathbb{S}^2} H_{10} d\sigma = \int_{\mathbb{S}^2} H_6 H_{10} d\sigma = 0, \quad \int_{\mathbb{S}^2} H_6^2 d\sigma = \frac{1584}{325}.$$

Lemma 3.1 (Invariant test set). *Let Q be an H_3 -invariant rule on $\mathbb{S}^2$. Then Q is exact for every spherical polynomial of degree at most t if and only if*

$$Q[H_6^r H_{10}^s] = \int_{\mathbb{S}^2} H_6^r H_{10}^s d\sigma \quad \text{whenever} \quad 6r + 10s \leq t.$$

Proof. The invariant ring of the reflection group H_3 has homogeneous basic degrees 2, 6, 10. On $\mathbb{S}^2$, the degree-two invariant is the constant 1. Replacing the degree-six and degree-ten generators by their harmonic parts is a triangular change of generators, so the restricted invariant algebra is generated by H_6 and H_{10} . Sobolev's invariant-cubature criterion then says that an H_3 -invariant rule is exact through degree t if and only if it is exact on all invariant polynomials through degree t [8]. The displayed monomials are a basis for those restricted invariants. $\square$

Every spherical moment used below is obtained from the exact monomial identity

$$\int_{\mathbb{S}^2} x^{2i} y^{2j} z^{2k} d\sigma = \frac{(2i-1)!!(2j-1)!!(2k-1)!!}{(2i+2j+2k+1)!!},$$

with $(-1)!! = 1$; a monomial with an odd exponent has mean zero. Consequently all moment matrices and right-hand sides lie in $\mathbb{Q}(\sqrt{5})$.

4 Exact positive rules for frequencies one through eight

Define

$$t_f = \begin{cases} 3f + 2, & f \text{ odd}, \\ 3f + 3, & f \text{ even}. \end{cases}$$

Theorem 4.1 (Certified fixed-grid cubature). *For every $1 \leq f \leq 8$, there is a unique H_3 -orbit-constant rule on X_f satisfying all invariant moment equations through degree t_f . Every orbit mass is strictly positive. The rule has exact algebraic degree t_f .*

f	$ X_f $	orbits	t_f	first failure	smallest W , approx.
1	12	1	5	6	1.000000
2	42	2	9	10	0.238095
3	92	3	11	12	0.080984
4	162	4	15	16	0.053869
5	252	5	17	18	0.030520
6	362	7	21	22	0.021749
7	492	8	23	24	0.015328
8	642	10	27	28	0.012087

Proof. For $\alpha = (a, b, c) \in \Lambda_f$, form the invariant moment vector

$$v_\alpha = (H_6(x_\alpha)^r H_{10}(x_\alpha)^s)_{6r+10s \leq t_f}$$

and the spherical target

$$b_f = \left(\int_{S^2} H_6^r H_{10}^s d\sigma \right)_{6r+10s \leq t_f}.$$

Let M_f have columns v_α . For $1 \leq f \leq 8$, the number of rows equals $|\Lambda_f|$. Exact inversion over $\mathbb{Q}(\sqrt{5})$ gives

$$M_f W_f = b_f.$$

The exact verifier constructs this inverse, reduces every entry to $a + b\sqrt{5}$ with $a, b \in \mathbb{Q}$, checks all residuals exactly, and proves every coordinate of W_f positive by rational sign comparisons. Lemma 3.1 gives exactness through t_f .

At the least invariant degree above t_f , the verifier evaluates, respectively for $f = 1, \dots, 8$,

$$H_6, H_{10}, H_6^2, H_6 H_{10}, H_6^3, H_6^2 H_{10}, H_6^4, H_6^3 H_{10}.$$

Every rule-minus-sphere residual is exactly nonzero. Thus the rule fails at that degree, proving that its algebraic degree is exactly t_f . $\square$

4.1 Compact exact formulas for $f \leq 4$

The first four rules fit legibly in the text. The orbit masses are

$$\begin{aligned} W_{100} &= 1, \\ W_{200} &= \frac{5}{21}, \quad W_{110} = \frac{16}{21}, \\ W_{300} &= \frac{26375 - 4500\sqrt{5}}{201432}, \\ W_{210} &= \frac{1570589 - 397404\sqrt{5}}{1007160}, \\ W_{111} &= \frac{-28971 + 17496\sqrt{5}}{41965}, \end{aligned}$$

and

$$\begin{aligned} W_{400} &= \frac{70025 - 12000\sqrt{5}}{801801}, \\ W_{310} &= \frac{3058238 - 84270\sqrt{5}}{8819811}, \\ W_{220} &= \frac{18302 - 270\sqrt{5}}{89089}, \\ W_{211} &= \frac{1059800 + 81000\sqrt{5}}{2939937}. \end{aligned}$$

The exact first-failure residuals are

$$\frac{132}{25}, \quad -\frac{4199}{5250}, \quad -\frac{14784}{35425}, \quad \frac{627(391241 - 123930\sqrt{5})}{374245000},$$

for $f = 1, 2, 3, 4$, respectively. The complete exact coefficients and failure residuals for $f = 5, 6, 7, 8$ are preserved in the certificate and CSV identified in Section 10; decimals in the table are expository only.

4.2 Exact sign decisions

For completeness, if $w = a + b\sqrt{5}$, the verifier never decides its sign numerically. When a and b have opposite signs, it compares a^2 and $5b^2$ in $\mathbb{Q}$, with the direction selected from the known rational signs of a and b . Thus all strict inequalities in Theorem 4.1 reduce to finite integer comparisons.

5 The exact OMEGA coefficient-field representation

The cubature calculation natively produces coefficients in $\mathbb{Q}(\sqrt{5})$. The following result provides an alternate exact representation of that field. It does not change the derivation in Sections 2–4.

Theorem 5.1 (Quadratic-field identification). *Let T be an indeterminate and let λ denote the residue class of the polynomial variable in*

$$K_\Omega = \mathbb{Q}(T)[\lambda]/(\lambda^2 - T\lambda - T^2).$$

Then K_Ω is a field and

$$r := \frac{2\lambda}{T} - 1$$

satisfies $r^2 = 5$. The map

$$\iota : \mathbb{Q}(T, \sqrt{5}) \longrightarrow K_\Omega, \quad \iota(\sqrt{5}) = r,$$

fixing $\mathbb{Q}(T)$, is an isomorphism. In particular,

$$\boxed{\iota(a + b\sqrt{5}) = (a - b) + \frac{2b}{T}\lambda}, \quad a, b \in \mathbb{Q}.$$

Proof. The discriminant of $X^2 - TX - T^2$ is $5T^2$, which is not a square in $\mathbb{Q}(T)$; hence the polynomial is irreducible and the quotient is a field. In the quotient,

$$\left(\frac{2\lambda}{T} - 1\right)^2 - 5 = \frac{4}{T^2}(\lambda^2 - T\lambda - T^2) = 0.$$

Thus $\sqrt{5} \mapsto r$ defines an embedding. Conversely,

$$\lambda = \frac{T(r + 1)}{2},$$

so r generates the quotient over $\mathbb{Q}(T)$, proving surjectivity. Expanding $a + br$ gives the displayed formula. $\square$

The frozen OMEGA specialization uses

$$\Omega_0 = \varphi + \pi, \quad \tau_\Omega = \frac{1}{\Omega_0}, \quad \lambda_\Omega = \frac{\varphi}{\Omega_0} = \varphi\tau_\Omega.$$

At $(T, \lambda) = (\tau_\Omega, \lambda_\Omega)$,

$$\frac{2\lambda}{T} - 1 = 2\varphi - 1 = \sqrt{5},$$

so the representation recovers the ordinary positive real square root.

Corollary 5.2 (Exact translation of every certified weight). *Every orbit mass in Theorem 4.1 has an exact two-channel OMEGA normal form. For example,*

$$W_{300} = \frac{26375 - 4500\sqrt{5}}{201432}$$

becomes

$$W_{300} = \frac{30875T - 9000\lambda}{201432T}.$$

All 40 certified orbit masses are exported in both forms by the exact certificate.

Remark 5.3 (Direction of dependence). The logical arrows are

$$\begin{aligned} \text{Class-I geometry} &\longrightarrow \text{invariant moment system} \\ &\longrightarrow \text{positive coefficient in } \mathbb{Q}(\sqrt{5}) \\ &\longrightarrow \text{OMEGA normal form.} \end{aligned}$$

There is no arrow from Ω_0 , T , or λ_Ω back to the icosahedral moment equations.

6 Finite 15-bit replay and the first disagreement

The same frozen contract declares

$$\begin{aligned} S &= 10^{15}, & Q &= 339949771344778, & q &= \frac{Q}{S}, \\ D &= 10^{60}, \\ L_{60} &= 339949771344778929576702809321279193551009905014259416689444, \\ U_{60} &= L_{60} + 1, \end{aligned}$$

and the strict enclosure

$$L := \frac{L_{60}}{D} < \lambda_\Omega < \frac{U_{60}}{D} =: U.$$

Only this last strict enclosure is assumed in the two results below.

Proposition 6.1 (Unsigned 15-bit conformance). *Conditional on $L < \lambda_\Omega < U$,*

$$\lfloor \lambda_\Omega d \rfloor = \left\lfloor \frac{Qd}{S} \right\rfloor \quad (0 \leq d < 2^{15}).$$

Proof. Exact integer division checks, for all 32,768 values,

$$\left\lfloor \frac{Qd}{S} \right\rfloor = \left\lfloor \frac{L_{60}d}{D} \right\rfloor = \left\lfloor \frac{U_{60}d}{D} \right\rfloor.$$

Every number strictly between the two endpoints has the same floor. The computation is finite, exhaustive, and uses arbitrary-precision integers. $\square$

The package also checks the implementation boundary

$$Q \cdot 32767 < 2^{64} - 1, \quad Q \cdot 65535 > 2^{64} - 1,$$

which explains why the declared unsigned state width is 15 bits when the product is required to fit in an unsigned 64-bit word.

Theorem 6.2 (First disagreement, with minimality). *Conditional on $L < \lambda_\Omega < U$, the least nonnegative integer d for which*

$$\lfloor \lambda_\Omega d \rfloor \neq \left\lfloor \frac{Qd}{S} \right\rfloor$$

is

$$\boxed{d_* = 42,189,065}.$$

At this value,

$$\left\lfloor \frac{Qd_*}{S} \right\rfloor = 14,342,162, \quad \lfloor \lambda_\Omega d_* \rfloor = 14,342,163.$$

Proof. Exact cross multiplication gives the strict chain

$$\frac{846547}{2490212} < \frac{Q}{S} < \frac{14342163}{42189065} < L < \lambda_\Omega < U < \frac{13495616}{39698853}.$$

Moreover,

$$14342163 \cdot 2490212 - 846547 \cdot 42189065 = 1$$

and

$$13495616 \cdot 42189065 - 14342163 \cdot 39698853 = 1.$$

Thus the middle fraction is a Farey neighbor of the fraction on each side. Any reduced rational strictly between two determinant-one neighbors has denominator at least the sum of their denominators. Therefore

$$\frac{14342163}{42189065}$$

has the smallest denominator among all rationals in $(Q/S, U)$: any other rational in that interval lies in one of the two adjacent Farey intervals and has denominator strictly larger than 42,189,065.

If a floor disagreement occurred at a smaller positive d , the integer $m = \lfloor Qd/S \rfloor + 1$ would give, after reduction if necessary, a rational $m/d \in (Q/S, \lambda_\Omega] \subset (Q/S, U)$ with denominator below d_* , a contradiction.

At d_* , the middle numerator 14,342,163 lies strictly between $(Q/S)d_*$ and Ld_* , while the upper endpoint still has floor 14,342,163. Hence the two displayed floors follow. $\square$

7 The all-frequency convex formulation

The verified frequencies suggest the following deliberately limited statement.

Conjecture 7.1 (Positive fixed-Class-I cubature). *For every $f \geq 1$, there are strictly positive H_3 -orbit masses on X_f whose rule is exact for every spherical polynomial of degree at most*

$$t_f = \begin{cases} 3f + 2, & f \text{ odd,} \\ 3f + 3, & f \text{ even.} \end{cases}$$

The conjecture makes no maximality claim. It is proved here only for $1 \leq f \leq 8$; it remains open here for $f \geq 9$.

Proposition 7.2 (Relative-interior formulation). *Fix f and t . Let v_α be the invariant moment vectors from the proof of Theorem 4.1, including the constant coordinate, and let b be the spherical target vector. Strictly positive orbit masses exact through degree t exist if and only if*

$$b \in \text{relint conv}\{v_\alpha : \alpha \in \Lambda_f\}.$$

Proof. A positive orbit-mass vector expresses b as a convex combination using every v_α with positive coefficient. Any supporting affine functional that is constant at b must then be constant at every v_α , so b is in the relative interior.

Conversely, let c be the average of all v_α . If b is in the relative interior, then for sufficiently small $\varepsilon > 0$,

$$y = \frac{b - \varepsilon c}{1 - \varepsilon}$$

still lies in the convex hull. Combining any convex representation of y with the uniform positive representation of c gives a representation of b in which every coefficient is positive. $\square$

This converts the all-frequency problem into an exact finite-dimensional convex-feasibility problem and its separating-hyperplane dual. Numerical linear programming may guide the search, but it is not a proof unless its output is promoted to exact moment and sign certificates.

8 Compatibility with the Papers I–V global object

The multi-sorted construction of [7] is written here as

$$\mathfrak{G} = (\mathcal{S}_{\text{SRSD}}, \mathcal{A}_\Omega, \mathcal{D}_{\text{quant}}, \mathcal{D}_{\text{rat}}, \mathcal{H}_{\text{ico}}, \mathcal{C}_{\text{fin}}; \mathcal{J}).$$

Its icosahedral sort already contains the node sets X_f , their equal-weight measures, and the radial limiting measure. Its arithmetic sort contains the distinguished OMEGA field. The present cubature data attach to those two sorts, and to the exact SRSD coordinate layer, without identifying the distinct measures or dynamics.

8.1 Same nodes, distinct measures

The radial branch of $\mathfrak{G}$ uses

$$\mu_f = \frac{1}{|X_f|} \sum_{x \in X_f} \delta_x.$$

The certified orbit masses instead define

$$\nu_f = \sum_{\alpha \in \Lambda_f} \frac{W_{f,\alpha}}{|\mathcal{O}_{f,\alpha}|} \sum_{x \in \mathcal{O}_{f,\alpha}} \delta_x, \quad 1 \leq f \leq 8.$$

Every $W_{f,\alpha}$ is strictly positive and their sum is one. Thus ν_f and μ_f have the same full support X_f , but generally assign different masses to its orbits.

Proposition 8.1 (Exact cancellation of the first radial invariants). *For the invariant harmonics of Section 3,*

$$\nu_f(H_6) = 0 \quad (2 \leq f \leq 8), \quad \nu_f(H_{10}) = 0 \quad (3 \leq f \leq 8).$$

The remaining low-frequency values are

f	$\nu_f(H_6)$	$\nu_f(H_{10})$
1	132/25	988/625
2	0	−4199/5250.

Proof. The normalized spherical means of H_6 and H_{10} are zero. Theorem 4.1 is exact through degree at least six for $f \geq 2$ and through degree at least ten for $f \geq 3$, which gives the asserted zeros. The two exceptional rows are exact evaluations reproduced by the interface verifier. $\square$

This is the sharp measure-theoretic distinction between the two papers. For example, the equal-weight frequency-two rule has $\mu_2(H_6) = 33/100$, whereas $\nu_2(H_6) = 0$. The radial asymptotic formula and the positive limiting value $\mu_{\text{rad}}(H_6) = C_0 > 0$ in [7] remain statements about μ_f and μ_{rad} ; they are not inherited by the reweighted measures ν_f . Orbit reweighting cancels the obstruction on the fixed nodes rather than altering the radial geometry.

8.2 Arithmetic specialization and canonical records

The global object writes

$$\tau_0 = \frac{1}{\Omega_0}, \quad \lambda_\Omega = \frac{\varphi}{\Omega_0}, \quad \lambda_\Omega^2 - \tau_0 \lambda_\Omega - \tau_0^2 = 0.$$

The generic quotient of Theorem 5.1 specializes by

$$T \mapsto \tau_0, \text{quad} \lambda \mapsto \lambda_\Omega.$$

This specialization does not collapse the coefficient field: τ_0 is transcendental, since algebraicity of τ_0 would make $1/\tau_0 - \varphi = \pi$ algebraic. Consequently the generic identity becomes the exact global-object identity

$$\sqrt{5} \mapsto \frac{2\lambda_\Omega}{\tau_0} - 1, \quad a + b\sqrt{5} \mapsto (a - b) + \frac{2b}{\tau_0} \lambda_\Omega.$$

Every certified orbit mass therefore lies in the arithmetic sort $\mathcal{A}_\Omega$, and the canonical symbolic record Can_Ω specified in [7] applies to it. The verifier in this package certifies the algebraic normal forms; it does not silently substitute a new implementation for that source's frozen byte serialization.

8.3 Exact transport through SRSD coordinates

For a positive scale vector $s \in (0, \infty)^3$, let E_s and D_s be the exact encoder and decoder of [7]. Explicitly,

$$E_s(y) = (q_s(y), r_s(y)), \quad D_s(d, \rho) = \text{diag}(s)d + \rho, \quad D_s E_s = \text{id}_{\mathbb{R}^3}.$$

Define

$$\widehat{\nu}_{f,s} = (E_s)_\# \nu_f.$$

Then every integrable observable Y satisfies

$$\int Y(D_s e) d\widehat{\nu}_{f,s}(e) = \int_{\mathbb{S}^2} Y(x) d\nu_f(x).$$

In particular, for every spherical polynomial p of degree at most t_f ,

$$\int p(D_s e) d\widehat{\nu}_{f,s}(e) = \int_{\mathbb{S}^2} p d\sigma.$$

This is lossless coordinate transport of an already proved rule. Dropping a residual channel, rounding, or retaining only an SRSD quotient is a different record map and is not covered by the identity.

Theorem 8.2 (Conservative cubature extension). *Let $\mathcal{C}_{\text{cub}}$ contain the forty orbit masses, the measures ν_f , their exact degrees, and their first-failure residuals for $1 \leq f \leq 8$. Adjoining this sort and the three displayed interfaces gives*

$$\mathfrak{G}^{\text{cub}} = (\mathfrak{G}, \mathcal{C}_{\text{cub}}; \mathcal{J} \cup \mathcal{J}_{\text{cub}}).$$

This extension exists and is conservative in the following concrete sense: projection onto $\mathfrak{G}$ recovers every original component and gluing law unchanged, while projection onto the Class-I geometry and $\mathcal{C}_{\text{cub}}$ recovers Theorem 4.1. No floor dynamics, finite-limb record, or SRSD transport statement is used to solve the cubature moment systems.

Proof. The exact certificate constructs $\mathcal{C}_{\text{cub}}$ over $\mathbb{Q}(\sqrt{5})$, proves positivity and the moment identities, and evaluates every first failure. The arithmetic interface is Theorem 5.1 followed by the specialization above. The SRSD interface follows from $D_s E_s = \text{id}$ and the definition of pushforward measure. These add a new sort and new maps whose codomains are existing sorts; they alter none of the data or equations already present in $\mathfrak{G}$. The two stated projections and the dependency claim follow directly. $\square$

The complete dependency boundary for this single integrated construction is summarized below.

Layer	Exact interface used	Not claimed
Radial	same X_f ; μ_f and ν_f distinguished	radial asymptotics for ν_f
OMEGA	exact normal form for every $W_{f,\alpha}$	OMEGA origin of geometry or weights
SRSD	lossless decoded moment transport	compression or quotient-only exactness
Floor systems	15-bit replay remains intact	role in moment equations or positivity
Finite records	the frozen canonical record accepts each field element	a replacement byte serialization
Cubature	exact rules and first failures for $f \leq 8$	all-frequency proof, maximality, novelty, or priority

9 Provenance and literature boundary

Invariant cubature and icosahedral orbit quadrature are classical subjects. McLaren [4], Goethals and Seidel [2], Konyaev [3], Ahrens and Beylkin [1], Nozaki and Sawa [8], and Popov [9] are among the relevant starting points. Popov, in particular, describes icosahedrally invariant spherical cubature through substantially higher orders using movable orbit parameters [9]. The present question is narrower: positive orbit-constant weights on the already fixed, radially projected Class-I node sets.

No priority conclusion is drawn here. The $f = 1$ icosahedral design and the $f = 2$ 42-node rule should be treated as classical or implicit unless a complete historical audit establishes otherwise. The exact fixed-node calculations for $f = 3, \dots, 8$ likewise require a specialist literature search before any novelty language is considered.

The radial construction and degree-six normalization are imported from [5]. The OMEGA contract, algebra, and enclosure are frozen from [6]; the live-site snapshot is provenance only and is not an executable dependency. The ambient multi-sorted notation, exact SRSD transport law, and the distinction between the equal-weight radial measure and other attached sorts are imported from the Papers I–V synthesis [7]. Its local PDF snapshot is frozen by hash; none of its unrelated dynamical or foundational claims is needed for the cubature proof.

10 Reproducibility

The reproducibility record contains the manuscript source, the frozen contract, the native baseline source, exact scripts, machine-readable coefficients, and human-readable certificates. The principal frozen artifacts are:

Artifact	SHA-256
certificates/cubature_f1_f8_certificate.txt	e37e9d1e5f0f3cc6093640e0f1460dabfabcf294e529489631c2f50980cb6443
data/orbit_masses_f1_f8.csv	e10d2a4278bf2ae497cb06127a239a0f1af7bf0cffad60311bf6c279918e7946
certificates/omega_embedding_certificate.txt	90abccbb31c5682c13412708b145e885522acb60d9ad90bd693b5219ba069306
certificates/omega_15bit_bridge_certificate.txt	4ef4248b51b69f9776fbeb375c3d99d793db153a7e3bea3d25175672d8e2f0ac
omega/OMEGA_NAUGHT_1_0.txt	ad71b4d5b9395ab47fae8e78decbbbeb56aee760c9b20833358b8940eab68f8b8
certificates/global_object_interface_certificate.txt	f1f82ab92e117e8bcd599440a843374d5376711ab5b0771fab4bfb3093d05c22
global_object/Global_Mathematical_Object_Papers_I-V_v1.0.pdf	cbe30136476c14602545eaab3b0984d005cce6626378ddf8c402e28363c90ed2

With Python 3 and SymPy 1.14.0, run

```
./run_checks.sh.
```

The first script reconstructs all moment systems and weights rather than trusting the stored CSV. It verifies harmonicity, exact spherical moments, orbit counts, matrix inversion, zero residuals, strict signs, first failures, and the OMEGA channel export. The second verifies the quotient-algebra identity and all 40 specializations. The third exhausts the 15-bit domain and checks the Farey minimality certificate. The fourth verifies the global-object interface: common support, probability normalization, exact obstruction cancellation, arithmetic specialization, and preservation of the source-PDF hash.

The paper compiles with three runs of `pdflatex`; three runs are used so the table of contents and all cross-references are settled.

11 Conclusion

The fixed radial Class-I grids support a concrete sequence of exact positive cubature rules through frequency eight. Their proof is native to icosahedral geometry and invariant moment theory. The OMEGA quotient algebra then gives every resulting $\mathbb{Q}(\sqrt{5})$ coefficient an exact alternate two-channel representation, while the frozen 15-bit contract supplies an independent deterministic replay test with a proved first disagreement.

Relative to the Papers I–V global object, the result is a conservative new cubature sort. It reweights the existing Class-I node sets, embeds every weight in the existing OMEGA arithmetic sort, and survives exact SRSD coordinate transport. The equal-weight radial measure, its asymptotics, and the rational and real floor dynamics retain their original meanings.

This is the intended integration: exact geometry first, exact representation second, and explicit dependency boundaries throughout.

References

- [1] C. Ahrens and G. Beylkin, Rotationally invariant quadratures for the sphere, *Proceedings of the Royal Society A* **465** (2009), 3103–3125. [doi:10.1098/rspa.2009.0104](https://doi.org/10.1098/rspa.2009.0104).
- [2] J. M. Goethals and J. J. Seidel, Cubature Formulae, Polytopes, and Spherical Designs, in *The Geometric Vein*, Springer, 1981, 203–218. [doi:10.1007/978-1-4612-5648-9_13](https://doi.org/10.1007/978-1-4612-5648-9_13).

- [3] S. I. Konyaev, Quadratures of Gaussian type for a sphere invariant under the icosahedral group with inversion, *Mathematical Notes* **25** (1979), 326–329.
- [4] A. D. McLaren, Optimal Numerical Integration on a Sphere, *Mathematics of Computation* **17** (1963), 361–383. [doi:10.1090/S0025-5718-1963-0159418-2](https://doi.org/10.1090/S0025-5718-1963-0159418-2).
- [5] M. Newman, *Radial Icosahedral Sampling and Its First Harmonic Obstruction: Exact degree-six activation and Euler–Maclaurin coefficients*, Zenodo, 2026. [doi:10.5281/zenodo.22033917](https://doi.org/10.5281/zenodo.22033917).
- [6] M. Newman, *OMEGA NAUGHT Ω_0* , Zenodo, 2026. [doi:10.5281/zenodo.22048540](https://doi.org/10.5281/zenodo.22048540).
- [7] M. A. Newman, *Global Mathematical Object Associated with Papers I–V: Exact resolution, Omega arithmetic, floor dynamics, and radial icosahedral harmonics*, version 1.0, 22 August 2026. [online PDF](#).
- [8] H. Nozaki and M. Sawa, Note on Cubature Formulae and Designs Obtained from Group Orbits, *Canadian Journal of Mathematics* **64** (2012), 1359–1377. [doi:10.4153/CJM-2011-069-5](https://doi.org/10.4153/CJM-2011-069-5).
- [9] A. S. Popov, Cubature formulas on a sphere invariant under the icosahedral rotation group, *Numerical Analysis and Applications* **1** (2008), 355–361. [doi:10.1134/S199542390804006X](https://doi.org/10.1134/S199542390804006X).